

The deconfining phase transition in $SU(N_c)$ gauge theories*

B. Lucini^{a†}, M. Teper^a, U. Wenger^{a‡}

^aTheoretical Physics, Oxford University,
1 Keble Road, Oxford OX1 3NP, United Kingdom

We report on our ongoing investigation of the deconfining phase transition in $SU(4)$ and $SU(6)$ gauge theories. We calculate the critical couplings while taking care to avoid the influence of a nearby bulk phase transition. We determine the latent heat of the phase transition and investigate the order and the strength of the transition at large N_c . We also report on our determination of the critical temperature expressed in units of the string tension in the large N_c limit.

1. INTRODUCTION

The large- N_c physics of QCD and $SU(N_c)$ gauge theories is of great theoretical and phenomenological interest. Lattice calculations of string tensions, glueball masses etc. [1] have confirmed that a smooth large- N_c limit is achieved by keeping fixed the 't Hooft coupling $g^2 N_c$ and that, as expected, the leading corrections are $O(1/N_c^2)$.

An interesting question concerns the order of the deconfining phase transition at $N_c \rightarrow \infty$. It has been argued [2] that the transition in $SU(3)$ may be accidentally first order, due to a cubic term in the effective potential, but in general is second order in $SU(N_c)$. Our ongoing lattice study [3], on which we report here, is addressing this question and thereby improving on earlier results [4,5].

We use the standard plaquette action on $L^3 \times L_t$ lattices with periodic boundary conditions. The deconfining phase transition is studied by keeping L_t fixed and varying $\beta = 2N_c/g^2$ so as to pass the temperature $T = 1/a(\beta)L_t$ through the transition.

In order to distinguish the confining and the deconfining phases we define the spatial average $\bar{l}_p$ of the Polyakov loop and correspondingly the spatial average $\bar{u}_p$ of the plaquette variable $\text{Tr}U_p$. The two order parameters so defined can be used to characterise the order of the phase transition.

For a first order transition, e.g., we should find tunneling between the phases near the critical temperature T_c and subsequently a double peak structure in the probability distributions of $\bar{l}_p$ and $\bar{u}_p$.

In order to determine the critical couplings $\beta_c(V)$ at finite volume $V = L^3$ we define a normalised Polyakov loop susceptibility

$$\frac{\chi_l}{V} = \langle |\bar{l}_p|^2 \rangle - \langle \bar{l}_p \rangle^2$$

and, for each volume V , obtain χ_l as a continuous function of β using standard reweighting techniques [6]. Then we determine β_c from the location at which the susceptibility has its maximum.

In complete analogy we can define the specific heat $C(\beta)$ from the average plaquette $\bar{u}_p$,

$$\frac{1}{\beta^2} C(\beta) = \frac{\partial}{\partial \beta} \langle \bar{u}_p \rangle = N_p \langle \bar{u}_p^2 \rangle - N_p \langle \bar{u}_p \rangle^2,$$

where $N_p = 6L^3L_t$ is the total number of plaquettes. The value of β where C has its maximum provides a different definition for a critical coupling which however should agree with the one from χ_l in the thermodynamic limit. Moreover we can use the finite size scaling behaviour of the peak value $C(\beta_c, V)$ as a criterion for the determination of the order of the transition. To be more

*Talk by U. Wenger.

†EU Marie Curie fellow.

‡PPARC SPG fellow.




Figure 1. Finite size scaling of $\beta_c(V)$ for $L_t = 5$ in SU(4).

precise, we expect the leading finite size scaling

$$\frac{C(\beta_c, V)}{V} = \begin{cases} a_1 L^{\rho-d} + a_2 L^{-d}, & \text{2nd order,} \\ a_1 + a_2 L^{-d}, & \text{1st order,} \end{cases}$$

where $\rho < 3$ is some combination of critical exponents and $d = 3$.

An important point we need to address is the influence of a bulk phase transition which can be close to the physical deconfining transition. Our choices of L_t for the different gauge groups ensure that the two transitions are well separated and can unambiguously be distinguished.

2. T_c IN SU(4)

In SU(4) we perform a finite size scaling study at $L_t = 5$ on $L = 12, 14, 16, 18, 20$ lattices with between 200 and 300 ksweeps at each β -value and lattice size. For β -values close to β_c we find clear tunneling transitions both in $\bar{l}_p$ and $\bar{u}_p$. The probability distributions of these quantities exhibit clear double peaking which is characteristic for a first order transition.

In fig. 1 we plot our values of $\beta_c(V)$ against the inverse spatial volume expressed in units of the temperature T together with the extrapolations to infinite volume. We note that if the transition is first order, as ours clearly is, then the leading

finite- V correction should be

$$\beta_c(V) = \beta_c(\infty) - \frac{h}{VT^3}, \quad (1)$$

where h is independent of the lattice action. We find that all our values of β_c are consistent with eq. (1) and we obtain $\beta_c(\infty) = 10.63709(72)$ in the $V \rightarrow \infty$ limit. Moreover the value of $h = 0.09 \pm 0.02$ we thus extract is close to the SU(3) value, $h \simeq 0.1$ [7]. This suggests that h depends only weakly on N_c and we will therefore use this value to extrapolate to $V = \infty$ in the SU(6) case where we do not perform an explicit finite volume study.

From $\beta_c(\infty)$ we obtain the critical temperature T_c in terms of the string tension σ [1],

$$\frac{T_c}{\sqrt{\sigma}} = 0.6024 \pm 0.0045 \quad \text{at } a = 1/5T_c. \quad (2)$$

Our preliminary $L_t = 6$ calculations give a value $\beta_c(\infty) = 10.780(10)$ which translates to $T_c/\sqrt{\sigma} = 0.597(8)$. This, taken together with the value in eq. (2), yields the extrapolated continuum value

$$\lim_{a \rightarrow 0} \frac{T_c}{\sqrt{\sigma}} = 0.584 \pm 0.030 \quad \text{in SU(4)}. \quad (3)$$

In fig. 2 we plot $C(\beta_c)/V$ against $1/V$ and note that the scaling is certainly consistent with a first order one. The intercept at $1/V = 0$ provides a measure of the latent heat, Δe , since

$$\lim_{V \rightarrow \infty} \frac{C(\beta_c, V)}{\beta_c^2 V} \propto (\langle \bar{u}_{p,c} \rangle - \langle \bar{u}_{p,d} \rangle)^2 = \Delta e^2,$$

where $\bar{u}_{p,c}$ and $\bar{u}_{p,d}$ are the average plaquette values at β_c in the confined and deconfined phases, respectively. We obtain $\Delta e(N_c = 4) = 0.00197(5)$ and observe that if we take the $L_t = 4$ SU(3) latent heat in [8] and naively scale it to $L_t = 5$, we obtain $\Delta e(N_c = 3) \simeq 0.0013$ which is substantially smaller than our above SU(4) value. This shows explicitly that the SU(4) transition is more strongly first order than the SU(3) one.

3. T_c IN SU(6)

In SU(6) we perform a study on a $16^3 \times 6$ lattice at values of β close to $\beta = 24.845$. In physical




Figure 2. Normalised specific heat $C(\beta_c, V)/V$ in SU(4) at $L_t = 5$ plotted against $1/V$.

units this is like a 14^3 lattice with $L_t = 5$ and thus we are confident from our SU(4) calculations to be close to the thermodynamic limit.

As before we observe well defined tunneling between confined and deconfined phases, characteristic of a first order phase transition. Using the reweighting technique we extract a critical value $\beta_c = 24.850(3)$ which we extrapolate to $V = \infty$ using eq. (1) with the corresponding value of $h = 0.09$ from SU(4). We obtain $\beta_c(\infty) = 24.855 \pm 0.003$ which translates into

$$\frac{T_c}{\sqrt{\sigma}} = 0.588 \pm 0.002 \quad \text{at } a = 1/6T_c \quad (4)$$

using the values of the SU(6) string tension in [9]. Preliminary $L_t = 5$ calculations provide an estimate $\beta_c(\infty) = 24.519(33)$ which translates to $T_c/\sqrt{\sigma} = 0.580(12)$. Taken together with the value in eq. (4) this gives the continuum value

$$\lim_{a \rightarrow 0} \frac{T_c}{\sqrt{\sigma}} = 0.605 \pm 0.026 \quad \text{in SU(6)}. \quad (5)$$

As for the strength of the first order transition, we note that an estimate of the latent heat, Δe , in SU(6) is larger than the one on a directly comparable $16^3 6$ SU(4) calculation indicating that the transition is certainly not weakening as $N_c \rightarrow \infty$.

4. CONCLUSIONS

We find that SU(4) gauge theory at $L_t = 5$ reveals a first order deconfining phase transition with a latent heat that is not particularly small. We also find evidence that the SU(6) transition at $L_t = 6$ is first order as well and certainly not weaker than the one in SU(4). We come to that conclusion by comparing the appropriately rescaled latent heat expressed in terms of T_c for the different gauge groups. In doing so we take care to avoid confusion with a nearby bulk phase transition which, on our lattices, can unambiguously be identified.

The N_c -dependence of the critical temperature when expressed in units of the string tension, $T_c/\sqrt{\sigma}$, appears to be weak. From a fit we obtain

$$\frac{T_c}{\sqrt{\sigma}} = 0.582(15) + \frac{0.43(13)}{N_c^2}. \quad (6)$$

Finally, we conclude that the $SU(N_c)$ transition at $N_c = \infty$ is first order and not particularly weak.

REFERENCES

1. B. Lucini and M. Teper, JHEP 0106 (2001) 050 (hep-lat/0103027).
2. R. Pisarski and M. Tytgat, hep-ph/9702340, R. Pisarski, hep-ph/0203271.
3. B. Lucini, M. Teper and U. Wenger, hep-lat/0206029.
4. M. Wingate and S. Ohta, Phys. Rev D63 (2001) 094502 (hep-lat/0006016).
5. R. Gavai, Nucl. Phys. B633 (2002) 127 (hep-lat/0203015).
6. A. M. Ferrenberg and R. H. Swendsen, Phys. Rev. Lett. 63, 1195 (1989); *ibid.* 61, 2635 (1988).
7. B. Beinlich, F. Karsch, E. Laermann and A. Peikert, Eur. Phys. J. C6 (1999) 133 (hep-lat/9707023).
8. N. Alves, B. Berg and S. Sanielevici, Nucl. Phys. B376 (1992) 218 (hep-lat/9107002).
9. L. Del Debbio, H. Panagopoulos, P. Rossi and E. Vicari, JHEP 0201 (2002) 009 (hep-th/0111090); Phys.Rev. D65 (2002) 021501 (hep-th/0106185).